

Indication Range






















Prof. Dr. Wolfram Jäger

CESA / VASA - New Surgical Technique for the Treatment of:


Pelvic Organ Prolaps

and

Urge Urinary Incontinence

POP-Q Stage	Vaginal Stump Prolaps	Uterine Prolaps	Cystocele after Hysterectomy	Cystocele	Therapy	Clinical Diagnosis
POP-Q Stage 0						Patient with Urge-Symptoms
POP-Q Stage I					DynaMesh®-CESA or DynaMesh®-VASA	Minor Urge Symptoms: Holding 3-10 min. Miction by day 10-15 times
POP-Q Stage II						Heavy Urge Symptoms: Holding < 3 min. Miction by day > 15 times
POP-Q Stage III						If Urge Symptoms are indicated: different characteristic
POP-Q Stage IV						No Urge Symptoms
Situation after CESA/VASA					DynaMesh®-CESA DynaMesh®-VASA DynaMesh®-CESA and DynaMesh®-VASA implants are especially designed for the surgical treatment of pelvic organ prolaps related Urge Urinary Incontinence according to the particular CESA (CErvice -SAcropexy) and VASA (VAgino-SACropexy) surgery techniques.	

In Case of Incontinence after CESA/VASA

Situation after SIS soft / SIS direct soft	 <p>In case of incontinence after CESA/VASA-implantation a tension free vaginal tape treatment is recommended 6 months postoperatively. DynaMesh®-SIS soft (retrosymphysary) or DynaMesh®-SIS direct soft (transobturatoric).</p>	<p>DynaMesh®-SIS soft DynaMesh®-SIS direct soft</p> <p>The DynaMesh®-SIS soft structures have been especially developed for the treatment of female stress urinary incontinence. DynaMesh®-SIS soft and DynaMesh®-SIS direct soft serve to reinforce connective tissue structures and ligaments.</p>
---	--	--